

Pion form factors from lattice QCD with exact chiral symmetry

Takashi Kaneko*

KEK Theory Center, High Energy Accelerator Research Organization (KEK), Ibaraki 305-0801, Japan

School of High Energy Accelerator Science, The Graduate University for Advanced Studies (Sokendai), Ibaraki 305-0801, Japan

We calculate pion vector and scalar form factors in two-flavor lattice QCD and study the chiral behavior of the vector and scalar radii $\langle r^2 \rangle_{V,S}$. For a direct comparison with chiral perturbation theory (ChPT), chiral symmetry is exactly preserved by employing the overlap quark action. We utilize the all-to-all quark propagator in order to calculate the scalar form factor including the contributions of disconnected diagrams. A detailed comparison with ChPT reveals that two-loop contributions are important to describe the chiral behavior of the radii in our region of the pion mass $M_\pi \gtrsim 290$ MeV. From chiral extrapolation based on two-loop ChPT, we obtain $\langle r^2 \rangle_V = 0.409(23)(37) \text{ fm}^2$ and $\langle r^2 \rangle_S = 0.617(79)(66) \text{ fm}^2$, which are consistent with phenomenological analyses.

*6th International Workshop on Chiral Dynamics, CD09
July 6-10, 2009
Bern, Switzerland*

*E-mail: takashi.kaneko@kek.jp

1. Introduction

A detailed study of the chiral behavior of pion observables is an important subject towards a deep understanding of the low-energy dynamics of QCD. The pion vector form factor $F_V(q^2)$ has been precisely measured by experiments and the charge radius $\langle r^2 \rangle_V$ can be extracted in a model-independent way. Although a detailed comparison of $\langle r^2 \rangle_V$ between chiral perturbation theory (ChPT) and non-perturbative determinations in lattice QCD is very interesting, the chiral behavior of $\langle r^2 \rangle_V$ is distorted by explicit chiral symmetry breaking with the conventional lattice quark actions.

The chiral behavior of the scalar form factor $F_S(q^2)$ is another interesting subject. The scalar radius $\langle r^2 \rangle_S$ provides a determination of the low-energy constant (LEC) l_4 , which should be compared with that from F_π , and $\langle r^2 \rangle_S$ has 6 times enhanced chiral logarithm compared to $\langle r^2 \rangle_V$. Since there are no experimental processes directly related to $F_S(q^2)$, its direct determination is possible only through lattice QCD. It is, however, computationally very demanding to evaluate contributions of disconnected diagrams to $F_S(q^2)$ with the conventional simulation methods.

In this article, we present our calculation of these pion form factors in two-flavor lattice QCD. Chiral symmetry is exactly preserved by employing the overlap quark action [1] for a direct comparison of $\langle r^2 \rangle_{V,S}$ with ChPT at two loops. We use the so-called all-to-all quark propagator [2] to calculate $F_S(q^2)$ including the disconnected contributions. We refer readers to Ref. [3] for more detailed description of this study.

2. Simulation method

We simulate QCD with two flavors of degenerate up and down quarks using the Iwasaki gauge action and the overlap quark action. We also introduce a topology fixing term [4] into the lattice action to substantially reduce the computational cost. Our numerical simulations are carried out on a $N_s^3 \times N_t = 16^3 \times 32$ lattice at a lattice spacing of $a = 0.1184(21)$ fm, which is fixed from the heavy quark potential. In the trivial topological sector $Q=0$, we simulate four values of the bare up and down quark masses $m_{ud} = 0.015, 0.025, 0.035$ and 0.050 , which cover a range of the pion mass $290 \lesssim M_\pi [\text{MeV}] \lesssim 520$. Statistics are 100 independent configurations at each m_{ud} . We also simulate nontrivial topological sectors $Q = -2$ and -4 to study effects of the fixed topology. Further details on our configuration generation are presented in Ref.[5].

The matrix element $\langle \pi(p') | \mathcal{O}_\Gamma(p' - p) | \pi(p) \rangle$ can be extracted from the three-point function

$$C_{\pi \mathcal{O}_\Gamma \pi}(\Delta t, \Delta t'; \mathbf{p}, \mathbf{p}') = \frac{1}{N_s^3 N_t} \sum_{\mathbf{x}, t} C_{\pi \mathcal{O}_\Gamma \pi}(\mathbf{x}, t; \Delta t, \Delta t'; \mathbf{p}, \mathbf{p}'), \quad (2.1)$$

$$C_{\pi \mathcal{O}_\Gamma \pi}(\mathbf{x}, t; \Delta t, \Delta t'; \mathbf{p}, \mathbf{p}') = \sum_{\mathbf{x}', \mathbf{x}''} \langle \mathcal{O}_\pi(\mathbf{x}', t + \Delta t + \Delta t') \mathcal{O}_\Gamma(\mathbf{x}'', t + \Delta t) \mathcal{O}_\pi(\mathbf{x}, t)^\dagger e^{-i\mathbf{p}'(\mathbf{x}' - \mathbf{x}'')} e^{-i\mathbf{p}(\mathbf{x}'' - \mathbf{x})} \rangle, \quad (2.2)$$

where $\mathcal{O}_\Gamma$ is the vector current V_μ or scalar operator S , and $\mathcal{O}_\pi^\dagger$ represents an interpolating field to create the physical pion state. In the conventional method, one calculates the so-called point-to-all quark propagator $S_F(\mathbf{x}', \mathbf{x})$, which flows from a fixed lattice site $\mathbf{x}$ to any site $\mathbf{x}'$, by solving

$$\sum_{\mathbf{x}'} D(\mathbf{y}, \mathbf{x}') S_F(\mathbf{x}', \mathbf{x}) = \delta_{\mathbf{y}, \mathbf{x}}, \quad (2.3)$$

where D is the Dirac operator. Then, $C_{\pi \mathcal{O}_\Gamma \pi}(\mathbf{x}, t; \Delta t, \Delta t'; \mathbf{p}, \mathbf{p}')$ can be calculated by connecting the point-to-all propagators as shown in Fig. 1. However, we need to solve Eq. (2.3) for each lattice site




Figure 1: Connected (left-most diagram) and disconnected (middle diagram) three point functions. The right-most diagram with the vacuum expectation value (VEV) $\langle \mathcal{O}_\Gamma \rangle$ contributes to $F_5(0)$. Each line shows quark propagator S_F . We use $S_F(x, x') = \gamma_5 S_F(x', x)^\dagger \gamma_5$ for the spectator quark.

x to evaluate the disconnected quark loop $D^{-1}(x, x)$ as well as to average $C_{\pi\mathcal{O}_\Gamma\pi}(\mathbf{x}, t; \Delta t, \Delta t'; \mathbf{p}, \mathbf{p}')$ over the pion source location $x = (\mathbf{x}, t)$ as in Eq. (2.1). This needs prohibitively large CPU cost.

This difficulty can be avoided by constructing the all-to-all quark propagator, which contains the quark propagating from any lattice site to any site, in an effective way. Along the strategy proposed in Ref. [2], we prepare 100 low-lying modes of D for each gauge configuration, and their contribution to the all-to-all quark propagator is calculated exactly as

$$(S_F)_{\text{low}}(x, y) = \sum_{k=1}^{100} \frac{1}{\lambda_k} u_k(x) u_k^\dagger(y), \quad (2.4)$$

where λ_k and u_k represent k -th smallest eigenvalue and corresponding eigenvector. It is expected that this low-mode contribution dominates low-energy dynamics of pions. Possibly small contribution from the remaining high-modes can be estimated stochastically by the so-called noise method [6], which is not computationally intensive. We refer to Ref. [3] for more technical details on our method to calculate pion correlators using the all-to-all propagator.

We calculate three-point functions, $C_{\pi V_4\pi}$ and $C_{\pi S\pi}$, as well as the two-point function

$$C_{\pi\pi}(\Delta t; \mathbf{p}) = \frac{1}{N_s^3 N_t} \sum_{\mathbf{x}'} \sum_{\mathbf{x}, t} \langle \mathcal{O}_\pi(\mathbf{x}', t + \Delta t) \mathcal{O}_\pi(\mathbf{x}, t)^\dagger e^{-i\mathbf{p}(\mathbf{x}' - \mathbf{x})} \rangle. \quad (2.5)$$

We take 33 choices for the spatial momentum $\mathbf{p}^{(i)}$ with $|\mathbf{p}^{(i)}| \leq 2$, which cover a region of the momentum transfer $-1.7 \lesssim q^2 [\text{GeV}^2] \leq 0$ for the three-point functions. Note that the spatial momentum is shown in units of $2\pi/L$ in this article.

3. Determination of pion form factors

For a precise lattice calculation of the form factors, it is advantageous to construct an appropriate ratio of correlators so that some uncertainties, such as renormalization factors, cancel at least partially in the ratio [7]. In this study, we calculate effective value of the vector form factor from

$$F_V(\Delta t, \Delta t'; q^2) = \frac{2M_\pi}{E_\pi(|\mathbf{p}|) + E_\pi(|\mathbf{p}'|)} \frac{R_V(\Delta t, \Delta t'; |\mathbf{p}|, |\mathbf{p}'|, q^2)}{R_V(\Delta t, \Delta t'; 0, 0, 0)}, \quad (3.1)$$

$$R_V(\Delta t, \Delta t'; |\mathbf{p}|, |\mathbf{p}'|, q^2) = \frac{1}{N_{|\mathbf{p}|, |\mathbf{p}'|} \text{ fixed } |\mathbf{p}|, |\mathbf{p}'|} \sum \frac{C_{\pi V_4\pi}(\Delta t, \Delta t'; \mathbf{p}, \mathbf{p}')}{C_{\pi\pi}(\Delta t; \mathbf{p}) C_{\pi\pi}(\Delta t'; \mathbf{p}')}, \quad (3.2)$$

where $(1/N_{|\mathbf{p}|, |\mathbf{p}'|}) \sum_{\text{fixed } |\mathbf{p}|, |\mathbf{p}'|}$ represents the average over momentum configurations $\{(\mathbf{p}, \mathbf{p}')\}$ corresponding to the given value of q^2 . As shown in the left panel of Fig. 2, we observe a very clear




Figure 2: Left panel : effective value $F_V(\Delta t, \Delta t'; q^2)$ at $m_{ud}=0.025$, which is around a quarter of the physical strange quark mass. Right panel : statistical fluctuation of $C_{\pi V_4 \pi}(\Delta t, \Delta t'; \mathbf{p}, \mathbf{p}')$ at $(|\mathbf{p}|, |\mathbf{p}'|) = (\sqrt{2}, 0)$ and $\Delta t = \Delta t' = 7$. Filled and open symbols are results with and without averaging over the source location $(\mathbf{x}, t)$.

signal of $F_V(\Delta t, \Delta t'; q^2)$ with the statistical accuracy of typically 3–5 %. We note that the use of the all-to-all propagator enables us to achieve this high accuracy by averaging pion correlators over the location of the source operator $(\mathbf{x}, t)$ in Eqs. (2.1) and (2.5). The right panel of Fig. 2 demonstrates the remarkable reduction of the statistical fluctuation of $C_{\pi V_4 \pi}$ by this averaging.

We determine the vector form factor $F_V(q^2)$ by a constant fit to the effective value $F_V(\Delta t, \Delta t'; q^2)$, and include the correction due to the finite lattice volume estimated in one-loop ChPT [8]. Although we do not observe any significant Q dependence of $F_V(q^2)$, the spread in $F_V(q^2)$ among $Q=0, -2$, and -4 is taken as a conservative estimate of the systematic uncertainty due to the fixed topology.

The scalar form factor normalized at a certain momentum transfer q_{ref}^2 can be calculated from

$$\frac{F_S(\Delta t, \Delta t'; q^2)}{F_S(\Delta t, \Delta t'; q_{\text{ref}}^2)} = \frac{R_S(\Delta t, \Delta t'; q^2)}{R_S(\Delta t, \Delta t'; q_{\text{ref}}^2)}, \quad (3.3)$$

$$R_S(\Delta t, \Delta t'; q^2) = \frac{1}{N_{|\mathbf{p}|, |\mathbf{p}'| \text{ fixed}}} \sum_{|\mathbf{p}|, |\mathbf{p}'|} \frac{C_{\pi S \pi}(\Delta t, \Delta t'; \mathbf{p}, \mathbf{p}')}{C_{\pi \pi}(\Delta t; \mathbf{p}) C_{\pi \pi}(\Delta t'; \mathbf{p}')}. \quad (3.4)$$




Figure 3: Left panel: effective value of normalized scalar form factor $F_S(\Delta t, \Delta t'; q^2)/F_S(\Delta t, \Delta t'; q_{\text{ref}}^2)$ at $m_{ud}=0.025$. Right panel: scalar form factor $F_S(q^2)$ with (filled symbols) and without (open symbols) the contributions of the disconnected diagrams. Both data are normalized by a common value $F_S(q_{\text{ref}}^2)$ including the disconnected contribution.

At $q^2=0$, $C_{\pi S\pi}$ has an additional contribution shown in Fig. 1 due to the VEV of the scalar operator S . The subtraction of this contribution leads to a relatively large uncertainty in $F_S(\Delta t, \Delta t'; q^2)$ at $q^2=0$ than at $q^2 \neq 0$, as seen in the left of Fig. 3. Although the Feynman-Hellmann theorem $2F_S(0) = \partial M_\pi^2 / \partial m_{ud}$ provides a better determination of $F_S(0)$, it is subject to systematic uncertainties of the chiral extrapolation of M_π^2 . In our simulation setup, $F_S(q^2)$ has the smallest relative error at the smallest nonzero value of $|q_{\text{ref}}^2|$ with $|\mathbf{q}_{\text{ref}}| = 1$. We therefore use $F_S(q^2)$ normalized at this q_{ref}^2 in the following analysis.

We determine $F_S(q^2)/F_S(q_{\text{ref}}^2)$ from the effective value $F_S(\Delta t, \Delta t'; q^2)/F_S(\Delta t, \Delta t'; q_{\text{ref}}^2)$ in a similar way to $F_V(q^2)$. The right panel of Fig. 3 compares $F_S(q^2)$ to that without the contribution of the disconnected diagrams. We observe a significant deviation between the two data, which implies the importance of the disconnected contributions in a precision study of $F_S(q^2)$.

4. Parametrization of q^2 dependence

In Fig. 4, we plot the vector form factor $F_V(q^2)$ and normalized scalar form factor $F_S(q^2)/F_S(q_{\text{ref}}^2)$ as a function of q^2 . We observe that $F_V(q^2)$ is close to the vector meson dominance (VMD) hypothesis $1/(1 - q^2/M_\rho^2)$ with the vector meson mass M_ρ measured at simulated m_{ud} . We then assume that the small deviation due to the higher poles or cuts can be approximated by a polynomial of q^2 . The q^2 dependence of $F_V(q^2)$ is therefore parametrized as

$$F_V(q^2) = \frac{1}{1 - q^2/M_\rho^2} + a_{V,1}q^2 + a_{V,2}(q^2)^2 + a_{V,3}(q^2)^3 = 1 + \frac{1}{6}\langle r^2 \rangle_V q^2 + c_V(q^2)^2 + \dots \quad (4.1)$$

in order to determine the charge radius $\langle r^2 \rangle_V$ and the curvature c_V . This form describes our data well as shown in Fig. 4. Results for $\langle r^2 \rangle_V$ and c_V do not change significantly if we remove the cubic term or if we add higher order terms into the parametrization form.

Due to the lack of the knowledge about the scalar resonances at the simulated quark masses, we use a generic quartic form

$$F_S(q^2) = F_S(0) \left(1 + \frac{1}{6}\langle r^2 \rangle_S q^2 + c_S(q^2)^2 + a_{S,3}(q^2)^3 + a_{S,4}(q^2)^4 \right) \quad (4.2)$$




Figure 4: Vector form factor $F_V(q^2)$ (left panel) and normalized scalar form factor $F_S(q^2)/F_S(q_{\text{ref}}^2)$ (right panel) as a function of q^2 . Solid lines show the fit curve and its error. We also plot the q^2 dependence of $F_V(q^2)$ expected from the VMD model by the dashed line.

to parametrize the q^2 dependence of $F_S(q^2)$. Our data are described by this form reasonably well as in Fig. 4. The result for the scalar radius $\langle r^2 \rangle_S$ is stable against the removal of the quartic term as well as inclusion of higher order terms. Such a stability is, however, not clear in the curvature c_S due to its large statistical uncertainty. We leave a precise determination of c_S for future studies, and only use results for $\langle r^2 \rangle_S$ in the following analysis.

ChPT can provide a more unambiguous parametrization of $F_{V,S}(q^2)$. Figure 5 shows contributions to $F_V(q^2)$ from each order $(q^2)^n$ of a Taylor expansion of Eq. (4.1). We observe that $O(q^6)$ and higher order contributions, which are NNNLO and higher in ChPT, become a small (a few %) correction below $|q^2| \lesssim (550\text{MeV})^2$. Our values of $|q^2|$ are, however, outside of this region due to the use of the simple periodic boundary condition for quark fields. We therefore do not use a parametrization of the q^2 dependence based on ChPT in this study.

We note that the simulated values of the pion mass squared M_π^2 are smaller than $(520\text{MeV})^2$. The $O(q^6)$ contribution to $F_V(q^2)$ is small if $|q^2|$ is smaller than this value. The quark mass dependence of our data of $\langle r^2 \rangle_{V,S}$ is expected to be described by NNLO ChPT.




Figure 5: Contributions in the q^2 expansion of $F_V(q^2) - 1$ at $m_{ud} = 0.025$. Thin solid, dashed and dot-dashed lines show $O(q^2)$, $O(q^4)$ and higher order contributions. The thick solid line is their total.

5. Chiral extrapolation

We first compare our lattice data of the radii $\langle r^2 \rangle_{V,S}$ with NLO ChPT formulae [9]

$$\langle r^2 \rangle_V = -\frac{1}{NF^2} (1 + 6Nl_6^r) - \frac{1}{NF^2} \ln \left[\frac{M_\pi^2}{\mu^2} \right], \quad \langle r^2 \rangle_S = \frac{1}{NF^2} \left(-\frac{13}{2} + 6Nl_4^r \right) - \frac{6}{NF^2} \ln \left[\frac{M_\pi^2}{\mu^2} \right], \quad (5.1)$$

where $N = (4\pi)^2$ and F is the decay constant in the chiral limit. We fix F to our estimate from our study of the pion decay constant [10]. The renormalization scale is set to $\mu = 4\pi F$. The NLO fits are not quite successful as seen in Fig. 6. While our data of $\langle r^2 \rangle_V$ are fitted well with $\chi^2/\text{d.o.f.} \sim 0.1$, the value extrapolated to the physical quark mass $0.364(4) \text{ fm}^2$ is significantly smaller than experiment [12]. On the other hand, the NLO formula for $\langle r^2 \rangle_S$ with the enhanced chiral log fails to reproduce our data and leads to large $\chi^2/\text{d.o.f.} \sim 9$.

We also note that NLO in ChPT is not sufficient to describe the quark mass dependence of the curvature c_V . Although c_V has a NLO term $1/(60NF^2M_\pi^2)$ coming from non-analytic NLO contributions to $F_V(q^2)$, it dominates c_V well below the physical pion mass and fairly near the chiral limit (see Fig. 7 below), where the non-analytic contributions to $F_V(q^2)$ are important to be consistent with the existence of the $\gamma \rightarrow \pi\pi$ branch cut at $q^2 \gtrsim 0$. Since c_V characterizes the $O(q^4)$ dependence of $F_V(q^2)$, NNLO contributions are essential to describe its quark mass dependence.

We therefore extend our analysis to NNLO ChPT. The NNLO contributions to the radii are [11, 14]




Figure 6: Chiral fit of $\langle r^2 \rangle_V$ (left panel) and $\langle r^2 \rangle_S$ (right panel) using NLO ChPT formulae. In the left panel, we also plot the experimental value $\langle r^2 \rangle_V = 0.437(16)$ fm² from an analysis based on $N_f = 2$ ChPT [11] (diamond) and $0.452(11)$ fm² quoted by Particle Data Group [12] (star). The diamond in the right panel represents $\langle r^2 \rangle_S = 0.61(4)$ fm² obtained from an indirect determination through $\pi\pi$ scattering [13].

$$\Delta \langle r^2 \rangle_V = \frac{1}{N^2 F^4} \left(\frac{13N}{192} - \frac{181}{48} + 6N^2 r_{V,r}^r \right) M_\pi^2 + \frac{1}{N^2 F^4} \left(\frac{19}{6} - 12N l_{1,2}^r \right) M_\pi^2 \ln \left[\frac{M_\pi^2}{\mu^2} \right], \quad (5.2)$$

$$\begin{aligned} \Delta \langle r^2 \rangle_S = & \frac{1}{N^2 F^4} \left(-\frac{23N}{192} + \frac{869}{108} + 88N l_{1,2}^r + 80N l_2^r + 5N l_3^r - 24N^2 l_3^r l_4^r + 6N^2 r_{S,r}^r \right) M_\pi^2 \\ & + \frac{1}{N^2 F^4} \left(-\frac{323}{36} + 124N l_{1,2}^r + 130N l_2^r \right) M_\pi^2 \ln \left[\frac{M_\pi^2}{\mu^2} \right] - \frac{65}{3N^2 F^4} M_\pi^2 \ln \left[\frac{M_\pi^2}{\mu^2} \right]^2. \end{aligned} \quad (5.3)$$

The NNLO expression of c_V is

$$\begin{aligned} c_V = & \frac{1}{60N F^2} \frac{1}{M_\pi^2} + \frac{1}{N^2 F^4} \left(\frac{N}{720} - \frac{8429}{25920} + \frac{N}{3} l_{1,2}^r + \frac{N}{6} l_6^r + N^2 r_{V,c}^r \right) \\ & + \frac{1}{N^2 F^4} \left(\frac{1}{108} + \frac{N}{3} l_{1,2}^r + \frac{N}{6} l_6^r \right) \ln \left[\frac{M_\pi^2}{\mu^2} \right] + \frac{1}{72N^2 F^4} \ln \left[\frac{M_\pi^2}{\mu^2} \right]^2. \end{aligned} \quad (5.4)$$

The analytic terms with $r_{\{V,S\},\{r,c\}}^r$ represent contributions from $O(p^6)$ chiral Lagrangian. The linear combination $l_1^r - l_2^r/2$ appearing commonly in $\langle r^2 \rangle_V$ and c_V is denoted by $l_{1,2}^r$ for simplicity.




Figure 7: Simultaneous chiral fit to $\langle r^2 \rangle_V$ and c_V based on two-loop formulae Eqs. (5.2) and (5.4). We also plot a phenomenological estimate $c_V = 3.85(60)$ [11].




Figure 8: Chiral extrapolation of radii $\langle r^2 \rangle_{V,S}$ obtained from simultaneous fit to $\langle r^2 \rangle_{V,S}$ and c_V .

We first carry out a simultaneous fit to $\langle r^2 \rangle_V$ and c_V in terms of $M_\pi^2/(NF^2)$. This fit has only four free parameters l_6^r , $l_{1,2}^r$, $r_{V,r}^r$ and $r_{V,c}^r$ [3], and these can be determined with reasonable accuracy without introducing phenomenological inputs. As seen in Fig. 7, our data are well described by the NNLO formulae with $\chi^2/\text{d.o.f.} \sim 0.7$. The extrapolated values of $\langle r^2 \rangle_V$ and c_V are consistent with recent phenomenological determinations [11, 15, 16].

The inclusion of $\langle r^2 \rangle_S$ into the simultaneous fit introduces additional four free parameters l_2^r , l_3^r , l_4^r and $r_{S,r}^r$, and we need to fix some of them to obtain a stable fit. In this study, we use a phenomenological estimate $\bar{l}_2 = 4.31(11)$ [13] and a lattice estimate $\bar{l}_3 = 3.38(56)$ from our analysis of the pion mass [10]¹, since they are determined with a reasonable accuracy and appear only in the NNLO terms. As plotted in Fig. 8, this fit describes our data of $\langle r^2 \rangle_S$ reasonably well with $\chi^2/\text{d.o.f.} \sim 0.7$. The extrapolations of $\langle r^2 \rangle_V$ and c_V are consistent with those in Fig. 7.

At physical quark mass, we obtain

$$\langle r^2 \rangle_V = 0.409(23)(37) \text{ fm}^2, \quad \langle r^2 \rangle_S = 0.617(79)(66) \text{ fm}^2, \quad c_V = 3.22(17)(36) \text{ GeV}^{-4}, \quad (5.5)$$

where the first and second errors are statistical and systematic, respectively. The latter includes uncertainties due to the choice of the input to fix the lattice scale and the inputs for the LECs (F , l_2 and l_3) as well as uncertainties due to the chiral extrapolation and lattice discretization. These results for $\langle r^2 \rangle_{V,S}$ and c_V are consistent with phenomenological analyses.

The results for the relevant LECs are

$$\bar{l}_6 = 11.9(0.7)(1.0), \quad \bar{l}_4 = 4.09(50)(52), \quad \bar{l}_1 - \bar{l}_2 = -2.9(0.9)(1.3), \quad (5.6)$$

$$r_{V,r}^r = -1.0(1.0)(2.5) \times 10^{-5}, \quad r_{V,c}^r = 4.00(17)(64) \times 10^{-5}, \quad r_{S,r}^r = 1.74(36)(78) \times 10^{-4}. \quad (5.7)$$

Our estimate of $\bar{l}_6$ is slightly smaller than those from ChPT analyses: $\bar{l}_6 = 16.0(0.9)$ from F_V [11] and $15.2(0.4)$ from τ and π decays [17]. This is partly because our estimate of F [10] is slightly smaller than phenomenological estimates. We note that $\bar{l}_4$ is consistent with our determination $\bar{l}_4 = 4.12(56)$ from F_π [10] and a phenomenological estimate $4.39(22)$ [13].

¹The μ -independent convention $\bar{l}_i$ is defined by $l_i^r = \gamma_i(\bar{l}_i + \ln[M_\pi^2/\mu^2])/2N$ with $\gamma_1 = 1/3$, $\gamma_2 = 2/3$, $\gamma_3 = -1/2$, $\gamma_4 = 2$ and $\gamma_6 = -1/3$.

6. Conclusions

In this article, we present our calculation of pion form factors in two-flavor lattice QCD with exact chiral symmetry, which enables us to unambiguously compare our lattice data with two-loop ChPT. By employing the all-to-all quark propagators, $F_S(q^2)$ is calculated including contributions from the disconnected diagrams for the first time. We observe that two-loop contributions are important to describe the quark mass dependence of $\langle r^2 \rangle_{V,S}$ and c_V at our region of the pion mass $M_\pi \gtrsim 290$ MeV. Our chiral extrapolation of $\langle r^2 \rangle_{V,S}$ and c_V are consistent with phenomenological analyses. We also confirm that $F_S(q^2)$ and F_π lead to consistent results for l_4 .

For a more precise comparison with experiment, we need to extend this study to three-flavor QCD. Such simulations are currently underway. Another important subject is a better control of the parametrization of the q^2 dependence of $F_{V,S}(q^2)$. To this end, the use of the twisted boundary condition [18] to simulate small values of $|q^2|$, dispersive analysis of the q^2 dependence [15, 16], and model-independent determination of the scalar resonance mass [19] are interesting possibilities for our future studies.

I am grateful to my colleagues in the JLQCD and TWQCD collaborations. I also thank Bala-subramanian Ananthanarayan and Sunethra Ramanan for a useful correspondence. Numerical simulations are performed on Hitachi SR11000 and IBM System Blue Gene Solution at High Energy Accelerator Research Organization (KEK) under a support of its Large Scale Simulation Program (No. 08-05). This work is supported in part by the Grant-in-Aid of the Ministry of Education (No. 20105005 and 21684013).

References

- [1] R. Narayanan and H. Neuberger, Nucl. Phys. B **443**, 305 (1995).
- [2] J. Foley *et al.* (TrinLat collaboration), Comput. Phys. Commun, **172**, 145 (2005).
- [3] S. Aoki *et al.* (JLQCD and TWQCD collaborations), Phys. Rev. D **80**, 034508 (2009).
- [4] H. Fukaya *et al.* (JLQCD collaboration), Phys. Rev. D **74**, 094505 (2006).
- [5] S. Aoki *et al.* (JLQCD collaboration), Phys. Rev. D **78**, 014508 (2008).
- [6] S.-J. Dong and K.-F. Liu, Phys. Lett. B **328**, 130 (1994).
- [7] S. Hashimoto *et al.*, Phys. Rev. D **61**, 014502 (1999).
- [8] T.B. Bunton, F.-J. Jiang and B.C. Tiburzi, Phys. Rev. D **74**, 034514 (2006).
- [9] J. Gasser and H. Leutwyler, Ann. Phys. **158**, 142 (1984).
- [10] J. Noaki *et al.* (JLQCD and TWQCD collaborations), Phys. Rev. Lett. **101**, 202004 (2008).
- [11] J. Bijnens, G. Colangelo and P. Talavera, JHEP **9805**, 014 (1998).
- [12] C. Amsler *et al.* (Particle Data Group), Phys. Lett. B **667**, 1 (2008).
- [13] G. Colangelo, J. Gasser and H. Leutwyler, Nucl. Phys. B **603**, 125 (2001).
- [14] J. Gasser and U.-G. Meißner, Nucl. Phys. B **357**, 90 (1991).
- [15] B. Ananthanarayan and S. Ramanan, Eur. Phys. J. C **60**, 73 (2009).
- [16] F.-K. Guo *et al.*, arXiv:0812.3270 [hep-ph].
- [17] M. González-Alonso, A. Pich and J. Prades, Phys. Rev. D **78**, 116012 (2008).
- [18] P.F. Bedaque, Phys. Lett. B **593**, 82 (2004).
- [19] I. Caprini, G. Colangelo and H. Leutwyler, Phys. Rev. Lett. **96**, 132001 (2006).